Competition Between Pressure and Gravity Confinement in Lyα Forest Observations

Jane C. Charlton[1,2], Edwin E. Salpeter[3] , and Suzanne M. Linder[1]

arXiv:astro-ph/9405005v1 3 May 1994

Abstract

A break in the distribution function of Lyα clouds (at a typical redshift of 2.5) has been reported by Petitjean et al. (1993). This feature is what would be expected from a transition between pressure confinement and gravity confinement (as predicted in Charlton, Salpeter, and Hogan (1993)). The column density at which the feature occurs has been used to determine the external confining pressure, $\sim 10\mathrm{cm}^{-3}\mathrm{K}$, which could be due to a hot, intergalactic medium. For models that provide a good fit to the data, the contribution of the gas in clouds to Ω is small. The specific shape of the distribution function at the transition (predicted by models to have a non-monotonic slope) can serve as a diagnostic of the distribution of dark matter around Lyα forest clouds, and the present data already eliminate certain models.

Subject headings: quasars: absorption lines - intergalactic medium - dark matter

Submitted to the *Astrophysical Journal Letters*

[1] Astronomy and Astrophysics Department, Davey Laboratory, Pennsylvania State University, University Park, PA 16802

[2] Center for Gravitational Physics and Geometry, Pennsylvania State University

[3] Space Sciences Building, Cornell Univ., Ithaca, NY 14853

1. Introduction

Clouds containing neutral hydrogen (with column densities ranging from $10^{13}\mathrm{cm}^{-2}$ up to $10^{22}\mathrm{cm}^{-2}$) are observed through Lyα absorption in quasar spectra. In this Letter we restrict ourselves to the epoch with the most reliable data, redshift $z \sim 2$ to 3. The distribution function $f(N_{Hobs})$ for observed HI column density of individual Lyα absorption lines is an important clue to the nature of the absorbing clouds, and the accuracy of determinations of f is improving steadily. With little loss of generality, an individual "Lyα forest cloud" with some N_{Hobs} can be modeled as a portion of a slab in hydrostatic equilibrium with perpendicular HI column density $N_H = N_{Hobs}\cos\theta$ (where θ is the inclination angle) and an (unknown) combined HI and HII column density N_{tot} (Charlton, Salpeter, & Hogan 1993 (CSH)). This modeling, in terms of a few adjustable parameters, is similar whether there are many different types of individual small clouds (minihalos), each with a different characteristic value of N_H, or a single type of very extended disk structure (large proto-galaxies) with N_H decreasing enormously from the center outwards.

Any realistic modeling of "forest clouds" must include the effects of both external pressure and self-gravity. CSH predicted a break in the distribution function, $f(N_{Hobs})$, where these two effects are equal. There must be a transition between pressure and gravity domination for some value N_{H1obs} of column density. The most recent observational data (see Petitjean et al. 1993 (PWRCL)) shows a break of the type predicted, with $N_{H1obs} \sim 10^{16}\mathrm{cm}^{-2}$. The transition value predicted by the CSH models depends on the value of the external pressure, P_{ext}, and weakly on the ratio of dark matter to gas density at the transition; thus we shall be able to estimate P_{ext}.

In this Letter we constrain model parameters by fitting to the neutral column density distribution over the forest regime. Since the observed distribution function $f(N_{Hobs})$ is of piece-wise power law form, it is a reasonable assumption that the basic distribution function g of the total (perpendicular) column density N_{tot} is a continuous power law. The observed $f(N_{Hobs})$ will be fit by adjusting three parameters: 1) the value of N_{H1} for the transition between pressure and gravity dominated regimes (set by the external pressure P_{ext} and a dark matter parameter η); 2) the slope of an assumed power law dependence of η on N_{tot} (The pressure induced by the self-gravity of the gas alone is proportional to N_{tot}^2, but the gravity due to dark matter present in the slab increases this pressure by a factor of η.) ; 3) the slope of the power law $g(N_{tot})$ distribution. We shall examine all of this three-dimensional parameter space, but give the specific example of extended proto-galactic disks in Section 3c.

Alternative models for the observed "break" have been given by PWRCL in terms of four types of clouds and a transition from metal-poor to metal-rich systems, but these require more parameters than the CSH model. We also omit, in this paper, possible deviations from ionization and thermal equilibrium (Miralda-Escudé and Rees 1993), since we wish to test the simplest models first.

Section 2 uses the relationship between neutral and total column density in various regimes (pressure dominated, gravity dominated, Lyman limit, and damped) to schematically illustrate the various features in the distribution function of N_H. Section 3 describes more detailed models that include corrections for orientation, extinction at the upper end of the forest regime, and consideration of the vertical slab structure. The best fit model

parameters (and plausible ranges) to the PWRCL column density distribution are presented. Also, the number of systems expected in the Lyman limit regime and the total contribution of forest systems to the mass density of the Universe are computed. Section 4 concludes with a summary of the constraints on the parameters. A value of P_{ext} is derived, and the implications of constraints on the dark matter distribution are discussed.

2. Overview of the Different Regimes

The distribution $g(N_{tot})$ is assumed to have a power law form. Written in logarithmic notation, $g'(N_{tot}) = N_{tot}g(N_{tot})$, and we define the parameter ϵ' by

$$g'(N_{tot})d\ln N_{tot} = CN_{tot}^{-\epsilon'}d\ln N_{tot}. \tag{1}$$

The distribution function $f'(N_H) = N_H f(N_H)$ for neutral hydrogen, in logarithmic notation, can be derived if N_H is given as a function of N_{tot}. Since $f'(N_H)d\ln N_H = g'(N_{tot})d\ln N_{tot}$ we have

$$f'(N_H) = g'(N_{tot})/k(N_{tot}),$$

$$k(N_{tot}) = d\ln N_H/d\ln N_{tot}. \tag{2}$$

To derive N_H as a function of N_{tot} in the forest regime, we model Lyα clouds as slabs in hydrostatic equilibrium and ionization/recombination balance. (For simplicity we neglect the presence of He.) The basic differences between behavior in the pressure and gravity dominated regimes can be illustrated by balancing the forces in a half slab approximation:

$$\frac{\pi}{2}G{m_H}^2\eta N_{tot}^2 + P_{ext} = 2n_{tot}kT \tag{3}$$

where n_{tot} represents the total number density of H (at a "typical" vertical distance from slab center). On the LHS, the first term is the force per unit area due to gravity, and P_{ext} is due to the inward, external force. The outward force (RHS) is due to gas pressure within the slab, with the factor of two arising because slabs are highly ionized so that free electrons also contribute to this pressure. The parameter η is the enhancement factor for gravity over the gas column due to dark matter, taken to depend on N_{tot} as

$$\eta(N_{tot}) = \eta_1(N_{tot}/N_{t1})^{-\delta} \tag{4}$$

where η_1 and N_{t1} are values at the transition between the pressure and gravity dominated regimes (where the two terms on the LHS of equation (3) are equal). For the case with no dark matter we have $\eta_1 = 1$ and $\delta = 0$. Using ionization/recombination balance for ionization rate ζ (obtained from the proximity effect (Bajtlik, Duncan, & Ostriker 1988), a temperature of 20000K, and equating the gravity and pressure terms in equation (3) gives P_{ext} in terms of the observable $N_{H1}(\sim N_{H1obs}/1.8)$ as

$$P_{ext}/k = 14.8\eta_1^{1/3}\left(\frac{\zeta}{2.7\times10^{-12}\mathrm{s}^{-1}}\times\frac{N_{H1obs}}{10^{16}\mathrm{cm}^{-2}}\right)^{2/3}\mathrm{cm}^{-3}\mathrm{K}. \tag{5}$$

The relationship between N_H and N_{tot} in terms of their values at the transition, N_{H1} and N_{t1}, is

$$\left(\frac{N_H}{N_{H1}}\right)_{hs} = 0.5\left[(N_{tot}/N_{t1})^{3-\delta} + (N_{tot}/N_{t1})\right] \tag{6}$$

where the subscript "hs" denotes the "half-slab" approximation used in this section.

Of necessity there are four different regimes for the relationship between N_H and N_{tot} in the range from the smallest to the largest values of column density, shown schematically in Figure 1a: As mentioned, there must be some value N_{t1} of N_{tot} (with a corresponding value N_{H1} of N_H) where the external and self-gravity pressures are equal. Far to the left of this transition (region (a) in Figure 1) the pressure is constant (= P_{ext}), N_H is proportional to N_{tot} (eq. (6)) and k (eq. (2)) is approximated by the constant $k_a = 1$. Far to the right of the transition (but still with $N_H << 10^{17.2}$) the ratio N_H/N_{tot} increases with increasing N_{tot} as $N_{tot}^{2-\delta}$, so that in this gravity dominated regime (b) the derivative k is approximated by the constant $k_b = 3 - \delta$. The next transition point (2) occurs when N_H is near the value $N_{H2} = 10^{17.2}\mathrm{cm}^{-2}$ where the optical depth for a (normally incident) photon just above the Lyman limit is 1. Just to the left of this transition, i.e. in the upper range of region (b), N_H/N_{tot} increases slightly more rapidly than $N_{tot}^{2-\delta}$ because photon extinction decreases the ionizing flux and hence increases the neutral fraction. To the right of this transition, over a fairly narrow range of N_{tot}, from $\sim N_{t2}$ to $N_{t3} \sim 2N_{t2}$, lies a region (c) which displays the "Stromgren effect", first suggested by Sunyaev (1969) and Felten and Bergeron (1969) and discussed in detail by Maloney (1993) and Corbelli and Salpeter (1993). The behavior of N_H in this region is complicated, but the essential point is that N_H increases by a very large factor, from N_{H2} to $N_{H3} \sim N_{t2} >> N_{H2}$. Finally, in the fourth region (d) where $N_{tot} >> N_{t3}$, the gas is almost entirely neutral, $N_H \approx N_{tot}$ and $k_d \approx 1$.

If equation (1) held for all values of N_{tot}, the distribution function $f'(N_H)$, given by equation (2), would also display the four regions shown in Figure 1b. If we define $\beta' = \beta - 1$ (where $f(N_H) \propto N_H^{-\beta}$) by

$$f'(N_H) \propto N_H^{-\beta'}, \tag{8}$$

then well into regions (a) and (d), $N_H \propto N_{tot}$ and $k_a = k_d = 1$ so that $\beta'_a = \beta'_d = \epsilon'$. Similarly, well into region (b), equation (8) holds with $\beta'_b = \epsilon'/k_b$ and $k_b = 3 - \delta$. However, just before the transition from a to b the function $k(N_{tot})$ increases before settling into the constant k_b, so that equation (2) predicts a short small dip, i.e. the slope $d\ln f'(N_H)/d\ln N_H$ does not decrease monotonically (see Figure 2a). The extinction correction (see section 3) near N_{H2} has a similar effect of increasing k and there is again an increase in slope. In the Stromgren effect region (c) the extinction increases rapidly and N_H/N_{tot} increases rapidly at first and then approaches unity, so that $k(N_H)$ increases at first, reaches a maximum and decreases towards unity. The actual behavior of $f'(N_H)$ in region (c), indicated schematically by the dashed curve in Figure 1b, depends on the shape of the photon spectrum and on just how large the ratio $N_{H3}/N_{H2} \sim N_{t2}/N_{H2} >> 1$ is (see Corbelli and Salpeter 1993), but $f'(N_H)$ is not much smaller at the high N_H end of region (c) than at the beginning. We shall see that $\epsilon' < 1$ for any acceptable model, so that the total gas mass would diverge on the upper end of N_{tot} if equation (1) held all the

way up. There must thus be another transition point (4) beyond which $g'(N_{tot})$ changes and eventually decreases rapidly in region (e).

The extent of the five regions (a) to (e) are not known a priori, nor are the transition values except for $N_{H2} \sim 10^{17.2}\mathrm{cm}^{-2}$ (defined by approximate unit optical depth at the Lyman edge). On the observational side there are three known transition points: $N_H \sim 10^{13}\mathrm{cm}^{-2}$ for the threshold of Lyα absorption observations, $N_H \sim N_{H2} \sim 10^{17.2}\mathrm{cm}^{-2}$ for the onset of the Lyman limit regime, and $N_H \sim 10^{20}\mathrm{cm}^{-2}$ for the onset of the damped wing regime. Between the last two points, numbers of absorption lines can be measured, but not the actual values of N_H and we shall refer to this range from $N_{Hobs} = 10^{17.68}\mathrm{cm}^{-2}$ to $N_{Hobs} = 10^{20.5}\mathrm{cm}^{-2}$ as "the long bin" of PWRCL. In principle, it is possible that $N_{H1} < 10^{13}\mathrm{cm}^{-2}$ so that all of the observable Lyα forest regime would lie in region (b), terminating at $N_{H2} \sim 10^{17.2}\mathrm{cm}^{-2}$. As Figure 1 shows, the predicted slope in this whole region (from $10^{13}\mathrm{cm}^{-2}$ to $10^{17.2}\mathrm{cm}^{-2}$) would have to be mostly constant with an increase at the end. The observational data of PWRCL suggests a decrease in slope, at best marginally compatible with a constant slope and incompatible with an increase in slope. We therefore accept their observational "break" near $N_{Hobs} \sim 10^{16}\mathrm{cm}^{-2}$ as representing the theoretically required change in slope at N_{H1} predicted by CSH.

3. Fitting More Accurate Models

We wish to fit all the observational data in the "Lyα forest regime" with $\log N_{Hobs}$ up to 17.68, using eq. (2) but adopting several model refinements.

a) Model Refinements

(*i*) Instead of the "half slab approximation" of section 2, we solve for the vertical structure of a slab, i.e. the density ρ as a function of height z (from slab center). We assume throughout that dark matter gravity dominates over that due to the gas, and that the dark matter density is approximately constant over the height of the gas slab, so that the gravitational acceleration $g(z)$ is proportional to z. With this assumption (plus assuming constant temperature) the shape of $\rho(z)$ can be calculated in terms of the parameter N_{tot}/N_{t1} alone (see CSH, Section 5.1).

(*ii*) In the approximate treatment so far, we have considered the ionization rate per H atom, ζ, a constant without allowance for absorption. By definition, observers chose for the upper bound to the "forest regime" (and lower limit to the "Lyman limit" regime, section 3d) a neutral column density N_{Hobs} where the optical depth is moderate, $\tau = 3(13.6\mathrm{eV}/E)^3$ for $10^{17.68}\mathrm{cm}^{-2}$). In the small perturbation approximation, the percentage increase in N_H/N_{tot} equals the decrease in ζ due to absorption, which depends only on $N_H = N_{Hobs}\cos\theta$ and the spectral shape (we use the usual quasar spectrum). We have calculated this small correction ($\lesssim 40\%$) for N_H up to $10^{17.41}$.

(*iii*) In reality, different slabs can have any inclination angle θ. The distribution of observed values of $f(N_{Hobs})$ can be derived from the distribution of perpendicular values (Milgrom 1988). Note that the orientation effect does not alter the power law behavior of $f(N_{Hobs})$ far from N_{H1} (on either side), but is responsible for a gradual transition instead of a sharp break.

b) Fits to N_H Distribution

For any choice of the model parameters N_{H1}, δ (eq. (4)), and ϵ' (eq. (1)), the distribution $f'(N_H)$ can be computed and normalized to provide the smallest value of χ^2 when compared to the PWRCL data. For various values of the dark matter parameter, δ, the best fit values of ϵ' and N_{H1} are given in Table 1. The smallest χ^2 value is obtained for $\delta \sim -1.5$, but the fit is only slightly worse for any $\delta \lesssim 0.65$. The slope of $f'(N_H)$ as a function of N_H is illustrated in Figure 2a for various δ, and the $f'(N_H)$ distributions are plotted along with the data in Figure 2b. Using $g'(N_{tot})$, which we have explicitly for the models in Table 1, the contribution to the cosmological density parameter is found to be $\Omega_{for} \sim 10^{-2}\eta_1^{-1/3}$ or slightly less for each of the four cases.

c) Extended Proto-Galaxy Disks

Hoffman et al. (1993) have proposed that the outer regions of proto-galaxy disks are responsible for the Lyman forest lines and that the total column density $N_{tot}(R)$ has a power law form. The exponent is related to ϵ' in eq. (1) by

$$N_{tot}(R) \sim R^{-p}, \qquad p = 2/\epsilon'. \tag{8}$$

If we assume a spherical dark matter halo which gives a flat rotation curve, we can derive a unique relation between the parameters δ and ϵ': The gravitational force per unit area in the GDR (the first term in eq. (3)) is proportional to $N_{tot}W/R^2$ where $W = N_{tot}/n_{tot}$ is the slab width. Using equation (8) we find

$$\delta = 1 - 0.5\epsilon', \tag{9}$$

Since ϵ' in Table 1 is close to 0.71 (p=2.8) this gives $\delta = 0.65$. If the rotation velocity declines slowly at large distances δ would be smaller.

For assumed values for the constant rotation velocity V_c and the radius R_{13} where N_H has dropped to $10^{13}\mathrm{cm}^{-2}$, one can calculate the transition dark matter parameter η_1. We find $\eta_1 = 4.7\ (V_c/70\mathrm{km/s})\ (R_{13}/250\mathrm{kpc})$ in terms of values for V_c and R_{13} suggested by the "Vanishing Cheshire Cat model" (Salpeter 1993; Hoffman et al. 1993). For these values $P_{ext}/k = 7\mathrm{cm}^{-3}\mathrm{K}$, and the Ly$\alpha$ forest systems come from radii between 24kpc and R_{13}, with a total gas mass of $8 \times 10^9 \mathrm{M}_\odot$ per proto-galaxy. The dark matter mass (out to R_{13}) is $3 \times 10^{11}\mathrm{M}_\odot$ per proto-galaxy, and the number density of these galaxies is about $4.2\mathrm{Mpc}^{-3}$ at $z = 2.5$ (the spheres would touch at $z \sim 3.3$). The contribution to Ω from the dark matter spheres is ~ 0.2.

d) Model Predictions for Lyman Limit Systems

In section 3b we obtained acceptable model fits to the observed distribution function $f'(N_{Hobs})$ up to $\log N_{Hobs} = 17.68$. In the "Lyman limit regime" we presently only have an observational value $I_{lb} = 0.42$ for the integral of f over a "long bin" stretching from $\log N_{Hobs} = 17.68$ to $\log N_{Hobs} = 20.5$ (PWRCL). Using equations (1) and (2) we calculate the integral of g' over the corresponding interval of N_{tot} and equate it to the observed I_{lb}.

For each model in Table 1 we give the dimensionless ratio $F_{lb} = I_{lb}\epsilon'/k(N_{t2})f'(N_{H2})$. If the power law in equation (1) holds for N_{tot} up to N_{t3}, then F_{lb} is given by

$$F_{lb} = 1 - (N_{t3}/N_{t2})^{-\epsilon'} . \tag{10}$$

The inequality $F_{lb} < 1$ (the requirement that the model predicts the observed number of Lyman limit systems for a finite value of N_{t3}) is violated only for the model with $\delta = 1.5$

IV. Discussion

This letter has focused on fitting equilibrium slab models of Lyα forest clouds to the neutral column density distribution of Petitjean et al. (1993) for clouds at $z \sim 2.5$. We find that the best fit is provided by models that have a transition between pressure and gravity confinement at neutral column densities of $N_{H1} \sim 10^{15}\mathrm{cm}^{-2}$. The best fit for this N_{H1} yields a χ^2 of 1.78 for $\epsilon' = .70$ and $\delta = -1.5$. However, fits are nearly as good for a larger range of parameters. Roughly, we have $14.8 < \log N_{H1} < 15.5$, $.6 < \epsilon' < .8$, and $\delta < 1$. In comparison, the best fit model in which the clouds are all gravity confined ($N_{H1} < 10^{13}\mathrm{cm}^{-2}$) has $\epsilon' = 1.3$, $\delta = .7$ and $\chi^2 = 3.5$. Thus we maintain that the data requires a transition from pressure to gravity confinement.

Since this transition is found to be at $N_{H1obs} \sim 2 \times 10^{15}\mathrm{cm}^{-2}$, equation (6) yields $P_{ext}/k = 5.1\eta_1^{1/3}\mathrm{cm}^{-3}\mathrm{K}$. This pressure could be due to a diffuse, universal, hot intergalactic medium. The measured value is smaller than limits placed by the COBE y parameter (distortion of the microwave background expected from Compton scattering by the IGM electrons) (Mather, et al. 1994). CSH favored a larger value of P_{ext} based on extrapolating a single power law $g'(N_{tot})$ to the damped wing systems, but region (e) in Figure 1 need not satisfy this power law.

Models with $\delta > 1$ give a bad fit to the Lyα forest systems and underpredict the number of Lyman limit systems ($F_{lb} > 1$). In any case, the simplest models for the dark matter distribution suggest $0 < \delta < 1$ and the proto-galaxy disk model (with constant rotation velocity and a spherical halo) gives $\delta \sim 0.6$. However, a disk galaxy with a declining rotation velocity and/or a compressed spheroid for the inner dark matter halo would have an appreciably smaller value for δ.

Although we have considered only simple models in this Letter, we hope that two features of our results are fairly universal: 1) The external pressure P_{ext} is likely to be large enough to lead to a pressure/gravity transition at $\log N_{H1} \sim 15$ and the neutral fraction of the hydrogen gas is likely to be large enough so that the gas in Lyα forest systems contributes rather little to the cosmological mass density. 2) Because of the form of equation (2), the slope of the distribution function $f'(N_H)$ should be a non-monotonic function (see Figure 2a). The present observational data in Figure 2b already seem to corroborate this feature.

We acknowledge support from NASA grant NAGW-3571 at Penn State and NSF grant AST 91-19475 at Cornell, the careful reading of the referee, Phil Maloney, and the Aspen Center for Physics.

Table 1: Summary of Model Results

δ	ϵ'	$\log N_{H1}$	χ^2	F_{lb}
-1.5	$.70^{+.11}_{-.10}$	$15.0^{+.4}_{-.2}$	1.78	.21
0	$.71^{+.09}_{-.08}$	$15.1^{+.4}_{-.2}$	1.95	.37
.65	.71	$15.2^{+.3}_{-.2}$	2.18	.57
1.5	$.69^{+.07}_{-.07}$	$15.4^{+.9}_{-.8}$	3.10	1.24

References

Bajtlik, S., Duncan, R. C., & Ostriker, J. P. 1988, ApJ, 327, 570

Charlton, J. C., Salpeter, E. E., & Hogan, C. J. 1993, ApJ, 402, 493 (CSH)

Corbelli, E., & Salpeter, E. E. 1993, ApJ, 419, 104

Felten, J. E., & Bergeron, J. 1969, Astrophysics Letters, 4, 155

Hoffman, G. L., Lu, N. Y., Salpeter, E. E., Farhat, B., Lamphier, C., & Roos, T. 1993, AJ, 106, 39

Maloney, P. 1993, ApJ, 414, 41

Mather, J. C., et al. 1994, ApJ, 420, 439

Milgrom, M. 1988, A&A. 202, L9

Miralda-Escudé, J., & Rees, M. J. 1993, MNRAS, 260, 617

Petitjean, P., Webb, J. K., Rauch, M., Carswell, R. F., & Lanzetta, K. 1993, MNRAS, 262, 499 (PWRCL)

Salpeter, E. E. 1993, AJ, 106, 1265

Sunyaev, R. A. 1969, SvA, 13, 5

Figure Captions

Figure 1: A schematic illustration of the regimes of behavior expected for Lyα clouds. The relationship between the total column density N_{tot} and the neutral column density N_H is given in the top half of the figure, and the lower portion sketches the resulting distribution function (the total neutral column density contained in systems with a given N_H). The five regimes are: a) pressure dominated, b) gravity dominated, c) "Stromgren sphere" transition between optically thin and optically thick clouds, d) optically thick regime in which clouds are almost neutral, and e) high column density cut-off.

Figure 2: The distribution function for the best fit models. The solid line represents the extended proto-galaxy disk model. a) The slope of the distribution function illustrates a feature in $f'(N_H)$ that is most pronounced for large negative values of δ. b) The distribution function is shown along with the data (1σ error bars) from PWRCL for the overall best fit model with $\delta = -1.5$, and for the proto-galaxy disk model.